



A Reliable Research Partner in Life Science and Medicine

# **Human CTLA4 Antibody Pair Set**

Catalog No. E-KAB-0709 Applications ELISA

**Synonyms** CD152;CTLA-4;CELIAC3;GRD4;GSE;IDDM12

## **Kit components & Storage**

Title	Specifications	Storage
Human CTLA4 Capture Antibody	1 vial, 100 μ g	Store at -20°C. Avoid freeze / thaw
		cycles.
Human CTLA4 Detection Antibody	1 vial, 50 μL	Store at -20°C. Avoid freeze / thaw
(Biotin)		cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0709)		
		Human CTLA4 Capture Antibody	Human CTLA4 Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Human CTLA4 protein	Recombinant Human CTLA4 protein	
Information	Swissprot	P16410		
Product details	Reactivity	Human	Human	
	Host	Goat	Goat	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%	
		glycerol, pH 7.4	protective protein, 50% glycerol, pH	
			7.4	
	Purify	Affinity purification	Affinity purification	
	Specificity	Detects Human CTLA4 in ELISAs.		

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>



A Reliable Research Partner in Life Science and Medicine

## **Applications**

Human CTLA4 Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4ug/mL	Human CTLA4 Capture	
Capture		Antibody	10
			8 15
			Optical Density
ELISA	1:1000-1:10000	Human CTLA4 Detection	)ydo 0.1.
Detection		Antibody (Biotin)	•
			0.01
			Human CTLA4 Concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

### **Background**

CTLA-4 (Cytotoxic T-Lymphocyte Antigen 4),also known as CD152,is a protein that plays an important regulatory role in the immune system. CTLA-4 is expressed on the surface of helper T cells and transmits an inhibitory signal to T cells. Mutations in the CTLA-4 gene have been associated with various autoimmune diseases, such as systemic lupus erythematosus, insulin-dependent diabetes mellitus, and other autoimmune diseases.

Toll-free: 1-888-852-8623 Tel: Web: <a href="https://www.elabscience.com">www.elabscience.com</a> Ema

Tel: 1-832-243-6086 Email: techsupport@elabscience.com

Fax: 1-832-243-6017