

Recombinant Human Cyclophilin C/PPIC Protein (His&Trx Tag)



Catalog Number:PKSH032316

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Peptidyl-Prolyl Cis-Trans Isomerase C;PPIase C;Cyclophilin C;Rotamase C;PPIC;CYPC
Species	Human
Expression Host	E.coli
Sequence	Lys31-Asp182
Accession	P45877
Calculated Molecular Weight	33.8 kDa
Observed molecular weight	34 kDa
Tag	N-Trx-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20°C.
Formulation	Supplied as a 0.2 μ m filtered solution of 20mM PB, 150mM NaCl, 10% Glycerol, pH 7.4.
Reconstitution	Not Applicable

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Cyclophilin C is an enzyme (EC 5.2.1.8) found in both prokaryotes and eukaryotes that interconverts the cis and trans isomers of peptide bonds with the amino acid proline. Proline has an unusually conformationally restrained peptide bond due to its cyclic structure with its side chain bonded to its secondary amine nitrogen. Most amino acids have a strong energetic preference for the trans peptide bond conformation due to steric hindrance, but prolines unusual structure stabilizes the cis form so that both isomers are populated under biologically relevant conditions. Proteins with prolyl isomerase activity include cyclophilin, FKBP, and parvulin, although larger proteins can also contain prolyl isomerase domains.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017