

Recombinant Mouse ICOS Ligand/ICOSL Protein (His Tag)

Catalog Number: PKSM041331

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse ICOS Ligand/ICOSL protein Glu47-Lys279, with an C-terminal His
Calculated MW	27.1 kDa
Observed MW	40-60 kDa
Accession	Q9JHJ8
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



Background

For Research Use Only

Mouse ICOS ligand(B7-H2) is an approximately transmembrane glycoprotein in the B7 family of immune regulatory molecules. B7-H2 is expressed on antigen presenting cells such as B cells, macrophages, monocytes, and dendritic cells. It binds to ICOS on activated T cells, leading to both positive and negative effects on immune responses including its own down-regulation. The B7-H2 interaction with ICOS is costimulatory for T cell proliferation as well as the development of B cells, plasma cells, follicular helper T cells and germinal centers. B7-H2 contributes to the development of allergic asthma by enhancing Th2 biased immune responses, limiting Th17 responses, and promoting eosinophilic infiltration into the lung. Its activation of ICOS on Treg limits pulmonary inflammation and airway hyperresponsiveness, promotes the development of inhalational tolerance, and impairs antitumor immunity. In the thyroid, B7-H2 is up-regulated on thyrocytes during inflammation and promotes their proliferation and production of thyroid hormones.

For Research Use Only

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017