

Recombinant Human FGF-2/FGFb Protein (aa 143-288)

Catalog Number: PKSH032439

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|----------------------|---|
| Species | Human |
| Source | E.coli-derived Human FGF-2;FGFb protein Pro143-Ser288 |
| Calculated MW | 16.3 kDa |
| Observed MW | 17 kDa |
| Accession | P09038-4 |
| Bio-activity | Measured in a cell proliferation assay using BALB/c 3T3 cells. The ED ₅₀ for this effect is 0.3-2.0 ng/ml. |


Properties

| | |
|-----------------------|---|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 0.01 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 100mM NaCl, 0.02% Tween 80, pH7.5. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 95 % as determined by reducing SDS-PAGE.



Measured in a cell proliferation assay using BALB/c 3T3 cells. The ED₅₀ for this effect is 0.3-2.0 ng/ml.

Background

For Research Use Only

FGF-basic is a members of the Fibroblast Growth Factors (FGFs) family. The family constitutes a large family of proteins involved in many aspects of development including cell proliferation; growth; and differentiation. They act on several cell types to regulate diverse physiologic functions including angiogenesis; cell growth; pattern formation; embryonic development; metabolic regulation; cell migration; neurotrophic effects; and tissue repair. FGF-basic is a non-glycosylated heparin binding growth factor that is expressed in the brain; pituitary; kidney; retina; bone; testis; adrenal gland liver; monocytes; epithelial cells and endothelial cells.