A Reliable Research Partner in Life Science and Medicine

Recombinant Human Activin RIIA/ACVR2A Protein (His Tag)

Catalog Number: PKSH033783

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Human

Source HEK293 Cells-derived Human Activin RIIA; ACVR2A protein Ala20-Pro134, with an C-

terminal His

Calculated MW 14.4 kDa
Observed MW 28-38 kDa
Accession P27037

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Concentration Subject to label value.

Endotoxin $< 1.0 \text{ EU} \text{ per } \mu\text{g} \text{ of the protein as determined by the LAL method.}$

Storage Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.


Shipping This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel

packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as a 0.2 μm filtered solution of 20mM PB, 300mM NaCl, 10% Glycerol, 5%

Trealose, pH7.4.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Activin Receptor Type-2A is a protein that in humans is encoded by the ACVR2A gene. ACVR2A is an activin type 2 receptor. This gene encodes activin A type II receptor. Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins; composed of a ligand-binding extracellular domain with cysteine-rich region; a transmembrane domain; and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding; resulting in phosphorylation of type I receptors by type II receptors. Type II receptors are considered to be constitutively active kinases.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017