

A Reliable Research Partner in Life Science and Medicine

## Recombinant Mouse ICAM-1/CD54 Protein(Fc Tag)

Catalog Number: PDMM100165

Note: Centrifuge before opening to ensure complete recovery of vial contents.

_					
	00	rri		177	'n
ш	es	U 1	124		ш

**Species** Mouse

Source HEK293 Cells-derived Mouse ICAM-1/CD54 protein Gln28-Asn485, with C-terminal

**Calculated MW** 75.2 kDa **Observed MW** 90 kDa P13597 Accession

**Bio-activity** Not validated for activity

#### **Properties**

**Purity** > 90% as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


Shipping This product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% **Formulation** 

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



SDS-PAGE analysis of Mouse ICAM-1/CD54 proteins, 2µg/lane of Recombinant Mouse ICAM-1/CD54 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 90 KD

## **Background**

For Research Use Only

Toll-free: 1-888-852-8623 Fax: 1-832-243-6017 Tel: 1-832-243-6086 Web: www.elabscience.com

Email: techsupport@elabscience.com

# Elabscience®

#### Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Intercellular adhesion molecule-1 (ICAM-1, or CD54) is a 90 kDa member of the immunoglobulin (Ig) superfamily and is critical for the firm arrest and transmigration of leukocytes out of blood vessels and into tissues. ICAM-1 is constitutively present on endothelial cells, but its expression is increased by proinflammatory cytokines. The endothelial expression of ICAM-1 is increased in atherosclerotic and transplant-associated atherosclerotic tissue and animal models of atherosclerosis. Additionally, ICAM-1 has been implicated in the progression of autoimmune diseases. ICAM-1 is a ligand for LFA-1(integrin). When activated, leukocytes bind to endothelial cells via ICAM-1/LFA-1 interaction and then transmigrate into tissues. Presence with heavy glycosylation and other structural characteristics, ICAM-1 possesses binding sites for some immune-associated ligands and serves as the binding site for entry of the major group of Human Rhinovirus (HRV) into various cell types. ICAM-1 also becomes known for its affinity for Plasmodium falciparum-infected erythrocytes (PFIE), providing more of a role in infectious disease. Previous studies have shown that ICAM-1 is involved in inflammatory reactions and that a defect in ICAM-1 gene inhibits allergic contact hypersensitivity.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com