

## Recombinant Human CD83/HB15 Protein (Fc Tag)

**Catalog Number:** PKSH032226

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human CD83;HB15 protein Thr 20-Ala143, with an C-terminal Fc
<b>Calculated MW</b>	41.2 kDa
<b>Observed MW</b>	50-60 kDa
<b>Accession</b>	Q01151
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 90 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
	Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 90 % as determined by reducing SDS-PAGE.

### Background

CD83 antigen is a single-pass type I membrane protein which contains one Ig-like V-type (immunoglobulin-like) domain. CD83 is expressed by activated lymphocytes; Langerhans cells and interdigitating reticulum cells. It contains one Ig-like V-type (immunoglobulin-like) domain. ; the soluble CD83 has the opposite effect and has an immune inhibitory capacity. Due to its immune inhibitory function; CD83 may play a significant role in antigen presentation or the cellular interactions that follow lymphocyte activation.

### For Research Use Only