



A Reliable Research Partner in Life Science and Medicine

POMC Polyclonal Antibody

catalog number: AN007010L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

T		4.0	
Desc	т	att o	n
DC31	7.0		

Reactivity Mouse

Immunogen Recombinant Human POMC protein expressed by E.coli

Host Rabbit Isotype IgG

Purification Antigen Affinity Purification

Conjugation Unconjugated

Buffer PBS with 0.05% proclin 300, 1% protective protein and 50% glycerol,pH7.4

Applications Recommended Dilution

WB 1:500-1:1000

Data



Western blot with Anti POMC Polyclonal antibody at dilution of 1:1000. Lane 1: Mouse pituitary tissue lysate.

Observed-MW:12 kDa Calculated-MW:29 kDa

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

This gene encodes a preproprotein that undergoes extensive, tissue-specific, post-translational processing via cleavage by subtilisin-like enzymes known as prohormone convertases. There are eight potential cleavage sites within the preproprotein and, depending on tissue type and the available convertases, processing may yield as many as ten biologically active peptides involved in diverse cellular functions. The encoded protein is synthesized mainly in corticotroph cells of the anterior pituitary where four cleavage sites are used; adrenocorticotrophin, essential for normal steroidogenesis and the maintenance of normal adrenal weight, and lipotropin beta are the major end products. In other tissues, including the hypothalamus, placenta, and epithelium, all cleavage sites may be used, giving rise to peptides with roles in pain and energy homeostasis, melanocyte stimulation, and immune modulation. These include several distinct melanotropins, lipotropins, and endorphins that are contained within the adrenocorticotrophin and beta-lipotropin peptides. The antimicrobial melanotropin alpha peptide exhibits antibacterial and antifungal activity. Mutations in this gene have been associated with early onset obesity, adrenal insufficiency, and red hair pigmentation. Alternatively spliced transcript variants encoding the same protein have been described.

For Research Use Only

Tel: 400-999-2100 Web: www.elabscience.cn Email:techsupport@elabscience.cn