

## Recombinant Mouse MME/CD10/Neprilysin protein (His Tag)

Catalog Number: PDEM100281

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Mouse
<b>Source</b>	E.coli-derived Mouse MME protein Met461-Trp750, with an N-terminal His
<b>Calculated MW</b>	31.8 kDa
<b>Observed MW</b>	32 kDa
<b>Accession</b>	Q61391
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 10 EU/mg of the protein as determined by the LAL method
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Reconstitution</b>	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

CD10, also known as neprilysin, common acute lymphoblastic leukemia antigen (CALLA), or neutral endopeptidase (NEP), is a zinc-dependent transmembrane metallo-endopeptidase. CD10 cleaves peptides at the N-terminal side of hydrophobic amino acid residues and deactivates a variety of signaling peptides. CD10 was shown to be one of the markers for a natural killer cell-restricted progenitor in fetal and adult tissues. Studies also suggest CD10 as a phenotypic marker to distinguish mature neutrophils from immature neutrophils in patients with inflammation. In addition, CD10+ GPR77+ carcinoma-associated fibroblasts (CAFs) promote tumor formation and chemoresistance.

### For Research Use Only