ZBTB3 Polyclonal Antibody

catalog number: E-AB-19552



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse

Immunogen Synthetic peptide of human ZBTB3

Host Rabbit IgG **Isotype**

Purification Antigen affinity purification


Unconjugated Conjugation


buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications	Recommended Dilution

WB 1:500-1:2000 IHC 1:35-1:200

Data






Western blot analysis of A549 A172 HT-29 HEPG2 and Hela Immunohistochemistry of paraffin-embedded Human liver cell lysates using ZBTB3 Polyclonal Antibody at dilution of cancer tissue using ZBTB3 Polyclonal Antibody at dilution 1:550

of 1:55(×200)

Observed-MV:Refer to figures

Calculated-MV:62 kDa



Immunohistochemistry of paraffin-embedded Human breast cancer tissue using ZBTB3 Polyclonal Antibody at dilution of 1:55(×200)

Preparation & Storage

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Storage

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

ZBTB3 Polyclonal Antibody

catalog number: E-AB-19552



The BTB (Broad-Complex, Tramtrack and Bric a brac) domain, also known as the POZ (POxvirus and Zinc finger) domain, is an N-terminal homodimerization domain that contains multiple copies of kelch repeats and/or C2H2-type zinc fingers. Proteins that contain BTB domains are thought to be involved in transcriptional regulation via control of chromatin structure and function. ZBTB3 (zinc finger and BTB domain containing 3) is a 574 amino acid protein that contains one BTB (POZ) domain and two C2H2-type zinc fingers. Localized to the nucleus, ZBTB3 is thought to play a role in transcriptional regulation events. The gene encoding ZBTB3 maps to human chromosome 11, which houses over 1,400 genes and comprises nearly 4% of the human genome.