



A Reliable Research Partner in Life Science and Medicine

Recombinant Human IL-28A protein(His Tag)

Catalog Number: PKSH034112

Note: Centrifuge before opening to ensure complete recovery of vial contents.

\mathbf{T}			4.
	A C	crin	tion

Species Human

Source E.coli-derived Human IL-28A protein Pro 27-Val 200, with an C-terminal His

Calculated MW20.6 kDaObserved MW20 kDaAccessionQ8IZJ0

Bio-activity Measure by its ability to protect HepG2 cells infected with encephalomyocarditis

(EMC) virus. The ED₅₀ for this effect is \leq 20 ng/mL.

Properties

Purity > 98 % as determined by reducing SDS-PAGE.

Endotoxin < 0.1 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS,pH 8.0.

Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Background

IL-28A (Interferon-λ2,IFN-λ2), IL-28B/IFN-λ3, and IL-29/IFN-λ1 are type III interferons which are distantly related to IL-10 family and type I IFN family cytokines. Mature human IL-28A is an approximately 22-25 kDa protein that shares 66% amino acid (aa) sequence identity with mouse and rat IL-28A and shows cross-species activity. It shares 96% and 70% aa sequence identity with human IL-28B and IL-29, respectively. IL-28A promotes the Th1 polarization of dendritic cells in the airway and inhibits Th2 and Th17 mediated inflammation. IL-28A additionally exhibits anti-tumor activity, in part by enhancing IL-12 dependent anti-tumor CTL responses in vivo. In contrast, it is up-regulated in invasive bladder cancer where it promotes tumor cell migration.