A Reliable Research Partner in Life Science and Medicine

# AF/LE Purified Anti-Human CD273 Antibody[24F.10C12]

Catalog Number: GF11750

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Reactivity Human

Immunogen Recombinant Human CD273 protein

**Host** Mouse

**Isotype** Mouse IgG2a, κ **Clone** 24F.10C12

**Purification** >98%, Protein A/G purified

Conjugation None (AF/LE)


Buffer Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the

LAL method.

## Applications Recommended Dilution

**FCM**  $2 \mu g/mL(0.5 \times 10^6 - 1 \times 10^6 \text{ cells})$ 

#### Data



HEK293T cells transfected with pcDNA3.1 plasmid encoding Human CD273 gene were stained with 0.2 μg AF/LE Purified Anti-Human CD273 Antibody[24F.10C12] (Right) and 0.2 μg Mouse IgG2a, κ Isotype Control (Left), followed by Elab

Fluor<sup>®</sup> 647-conjugated Goat Anti-Mouse IgG Secondary Antibody.

# Preparation & Storage

Storage Storage Storage, avoid

freeze / thaw cycles. This preparation contains no preservatives, thus it

should be handled under as eptic conditions.

Shipping lce bag

# **Background**

T cells require a signal induced by the engagement of the T cell receptor and a "co-stimulatory" signal(s) through distinct T cell surface molecules for optimal T cell activation and tolerance. Members of the B7 superfamily of counter-receptors were identified by their ability to interact with co-stimulatory molecules found on the surface of T cells. Members of the B7 superfamily include B7-1 (CD80), B7-2 (CD86), B7-H1 (PD-L1), B7-H2 (B7RP-1), B7-H3, and PD-L2 (B7-DC).