

Recombinant Human BAI3 Protein (His Tag)

Catalog Number: PKSH032129

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human BAI3 protein Ala25-Thr880, with an C-terminal His
Calculated MW	97.5 kDa
Observed MW	115-145 kDa
Accession	O60242
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, 5% Thehalose, pH 7.2. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Human Brain-Specific Angiogenesis Inhibitor 3 (BAI3) is a 177 kDa seven-span transmembrane (TM) protein, which is thought to be a member of the secretin receptor family. It is synthesized by neurons of the CNS and likely is a negative regulator of angiogenesis. BAI3 is 1498 amino acids in size. It contains three distinct regions: an N-terminal extracellular domain (ECD) (aa25-883), a 7-TM segment, and a C-terminal cytoplasmic region. The ECD contains four antiangiogenic TSP type 1 repeat (aa296-508), and one GSP domain (aa 816-867) that is likely used to cleave the ECD from the membrane-bound receptor. There is one alternate splice form that shows a deletion of aa 643-665. Over aa 25-880, human BAI3 shares 98% aa identity with mouse BAI3. BAI3 has been reported primarily in the brain, but is also localized to lung, testis, and pancreas. It might be involved in angiogenesis inhibition and suppression of glioblastoma.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017