Recombinant Rat Cyt-C protein (GST Tag)


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Catalog Number: PDER100180



Description **Species** Rat 11.4 kDa Mol Mass Accession P62898 **Bio-activity** Not validated for activity **Properties** > 90% as determined by reducing SDS-PAGE. Purity Endotoxin < 10 EU/mg of the protein as determined by the LAL method Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 Storage °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at $< -20^{\circ}$ C for 3 months. This product is provided as lyophilized powder which is shipped with ice packs. Shipping Formulation Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol. Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Rat Cyt-C proteins, 2µg/lane of Recombinant Rat Cyt-C proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 28 KD.

Background

Electron carrier protein. The oxidized form of the cytochrome c heme group can accept an electron from the heme group of the cytochrome c1 subunit of cytochrome reductase. Cytochrome c then transfers this electron to the cytochrome oxidase complex, the final protein carrier in the mitochondrial electron-transport chain. Plays a role in apoptosis. Suppression of the anti-apoptotic members or activation of the pro-apoptotic members of the Bcl-2 family leads to altered mitochondrial membrane permeability resulting in release of cytochrome c into the cytosol. Binding of cytochrome c to Apaf-1 triggers the activation of caspase-9, which then accelerates apoptosis by activating other caspases.

For Research Use Only