# Purified Anti-Human CD40 Antibody[G28.5]

catalog number: E-AB-F1214A



Note: Centrifuge before opening to ensure complete recovery of vial contents.

## Description

**Reactivity** Human

Immunogen Recombinant Human CD40 protein

**Host** Mouse

**Isotype** Mouse IgGl, κ

Clone G28.5

**Purification** >98%, Protein A/G purified


**Conjugation** Unconjugated

buffer PBS, pH 7.2. Contains 0.05% proclin 300.

# Applications Recommended Dilution

FCM  $2 \mu g/mL(1\times10^5-5\times10^5 \text{ cells})$ 

#### Data



Human peripheral blood lymphocytes were stained with 0.2 μg AF/LE Purified Anti-Human CD40 Antibody[G28.5] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control(Left),

followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD19 PE-conjugated Monoclonal Antibody.

# **Preparation & Storage**

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

## Background

CD40 is a 48 kD type I glycoprotein also known as BP50. It is a member of the TNFR superfamily primarily expressed on B cells, macrophages, follicular dendritic cells, endothelial cells, fibroblasts, and at low levels on plasma cells. CD40 has been reported to be involved in B cell differentiation, costimulation, isotype class-switching, and protection of B cells from apoptosis. Additionally, CD40 is important for T cell-B cell interactions. The ligand of CD40 is CD154 (CD40 ligand)

For Research Use Only