

## Purified Anti-Mouse CD274/PD-L1 Antibody[10F.9G2]

Catalog Number: GF1132A

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description


Reactivity	Mouse
Immunogen	Recombinant Mouse CD274/PD-L1 protein
Host	Rat
Isotype	Rat IgG2b, $\kappa$
Clone	10F.9G2
Purification	>98%, Protein A/G purified
Conjugation	Unconjugated
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

### Applications

### Recommended Dilution

FCM	2 $\mu\text{g/mL}$ ( $0.5 \times 10^6$ - $1 \times 10^6$ cells)
-----	---

### Data



C57/BL6 Mouse splenocytes were stained with 0.2  $\mu\text{g}$  Purified Anti-Mouse CD274/PD-L1 Antibody[10F.9G2](Right) and 0.2  $\mu\text{g}$  Rat IgG2b,  $\kappa$  Isotype Control(Left), followed by Elab Fluor® 647-conjugated Goat Anti-Rat IgG Secondary Antibody.

### Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

### Background

CD274, also known as B7-H1 or programmed death ligand 1 (PD-L1), is a 40 kD type I transmembrane protein and a member of the B7 family within the immunoglobulin receptor superfamily. It is expressed on T cells, B cells, NK cells, dendritic cells, IFN- $\gamma$  activated endothelial cells, and monocytes. B7-H1 is one of the ligands of PD-1. The interaction of B7-H1 with PD-1 plays an important role in the inhibition of T cell responses. Other studies have shown that B7-H1 is able to costimulate T cell growth and cytokine production. CD274 is involved in costimulation essential for T cell proliferation and production of IL-10 and IFN- $\gamma$ , in an IL-2-dependent and a PD-1-independent manner. Its interaction with PD-1 inhibits T cell proliferation and cytokine production.