

LIF, Mouse, Recombinant**Cat. No. : GPCK307****产品信息**

物种	Mouse
表达宿主	E.coli
序列信息	SPLPITPVNATCAIRHPCHGNLMNQIKNQLAQLNGSANALFISYYTAQGEPFPNNVEKLCAPN MTDFPSFHGNGTEKTKLVELYRMVAYLSASLTNITRDQKVLNPTAVSLQVKLNATIDVMRGLL SNVLCRLCNKYRVGHVDVPPVPDHSDKEAFQRKKLGCGLLGTYKQVISVVVQAF with polyhistidine tag at the N-terminus.
检索号	P09056.1
标签	His-tag at the N-terminus
分子量	20.68 kDa
有效期	12 months
生物活性	Measure by its ability to induce IL-6 secretion in M1 cells. The ED50 for this effect is < 0.5 ng/mL. The specific activity of recombinant mouse LIF is > 2 × 10 ⁶ IU/mg.

产品特性

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 μm filtered solution containing 1 × PBS, pH 7.4.
复溶	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

LIF, a pleiotrophic factor, is identified in multiple cell types, including T cells, myelomonocytic lineages, fibroblasts, liver, heart and melanoma. LIF is capable of promoting long-term maintenance of embryonic stem cells by inhibiting spontaneous differentiation. In addition, LIF also have abilities including stimulation of differentiation of cholinergic nerves, the stimulation of acute phase protein synthesis by hepatocytes, and suppression of adipogenesis by supressing the lipoprotein lipase in adipocytes.

