

Recombinant Human EpCAM/TROP-1 Protein (Fc Tag)

Catalog Number: PKSH032383

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human EpCAM;TROP-1 protein Gln24-Lys265, with an C-terminal Fc
Calculated MW	54.5 kDa
Observed MW	60-80 kDa
Accession	AAH14785.1
Bio-activity	Immobilized Recombinant Human EpCAM (C-Fc)(PKSH032383) at 1µg/ml (100 µl/well) can bind Anti-Human EpCAM Antibody(Human IgG4): Biotinylated by NHS-biotin prior to testing.The ED ₅₀ of Anti-Human EpCAM Antibody is 13.82ng/ml.


Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.



Immobilized Recombinant Human EpCAM (C-Fc) (PKSH032383) at 1µg/ml (100 µl/well) can bind Anti-Human EpCAM Antibody(Human IgG4): Biotinylated by NHS-biotin prior to testing.The ED₅₀ of Anti-Human EpCAM Antibody is 13.82ng/ml.

Background

For Research Use Only

Epithelial Cell Adhesion Molecule (EpCAM) is a signal type I transmembrane glycoprotein that belongs to the EPCAM family. EpCAM is composed of an extracellular domain with one thyroglobulin type-1 domain; a transmembrane domain and a cytoplasmic domain. EpCAM is found on the surface of adenocarcinoma; but not on mesodermal or neural cell membranes. The EpCAM molecule has been shown to function as a homophilic Ca²⁺ independent adhesion molecule. It may act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium as an immunological barrier providing the first line of defense against infection. Defects in EPCAM are a cause of hereditary non-polyposis colorectal cancer type 8 (HNPCC8) and diarrhea type 5 (DIAR5). EpCAM plays a role in embryonic stem cells proliferation and differentiation; it up-regulates the expression of FABP5; MYC and Cyclin A and Cyclin E. It is highly and selectively expressed by undifferentiated embryonic stem cells.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017