

A Reliable Research Partner in Life Science and Medicine

Recombinant Mouse CD30 Ligand/TNFSF8 Protein(Fc Tag)

Catalog Number: PDMM100135

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Mouse

Source HEK293 Cells-derived Mouse CD30 Ligand/TNFSF8 protein Gln68-Asp239, with an

C-terminal Fc

Calculated MW 43.8 kDa 55-65 kDa **Observed MW** P32972 Accession

Bio-activity Not validated for activity

Properties

Purity > 90% as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


Shipping This product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% **Formulation**

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Mouse CD30 Ligand/TNFSF8 proteins, 2µg/lane of Recombinant Mouse CD30 Ligand/TNFSF8 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 55-65 KD

Background

Toll-free: 1-888-852-8623 Fax: 1-832-243-6017 Tel: 1-832-243-6086

Web: www.elabscience.com Email: techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

CD30 ligand (CD30L), also known as CD153 and TNFSF8, is a membrane-associated glycoprotein belonging to the TNF superfamily and TNFR superfamily, and is a specific ligand for CD30/TNFRSF8 originally described as a cell surface antigen and a marker for Hodgkin lymphoma and related hematologic malignancies. CD30L is a type-II membrane glycoprotein expressed on activated T cells, stimulated monocyte-macrophages, granulocytes, eosinophils, and some Burkitt-like lymphoma cell lines. CD30L is capable of transducing signals through CD30 on different CD30+ lymphoma cell lines, and mediates pleiotropic biologic effects including cell proliferation, activation, differentiation, as well as cell death by apoptosis. CD30-CD30 ligand interaction has been suggested to have a pathophysiologic role in malignant lymphomas, particularly Hodgkin disease, large cell anaplastic lymphomas and Burkitt lymphomas, and is also involved in activation and functioning of the T cell-dependent immune response. Thus, CD153 and its receptor CD30 are regarded as therapeutic targets in hematologic malignancies, autoimmune and inflammatory diseases.全称

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com