A Reliable Research Partner in Life Science and Medicine

Recombinant Human PD-L2/CD273 Protein (mFc Tag)

Catalog Number: PKSH033556

Note: Centrifuge before opening to ensure complete recovery of vial contents.

\mathbf{r}		crip					
	00	0	РΤ	n	ŤΤ		m
v	\mathbf{c}	v.		w	w	w	ш

Species Human

Source HEK293 Cells-derived Human PD-L2/CD273 protein Leu20-Pro219, with an C-terminal

mFc

Calculated MW 49.2 kDa
Observed MW 60-90 kDa
Accession Q9BQ51

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Programmed Cell Death 1 Ligand 2 (PDCD1LG2) is a member of the BTN/MOG family. PDCD1LG2 contains one Ig-like C 2-type domain and one Ig-like V-type domain. PDCD1LG2 is highly expressed in the heart; placenta; pancreas; lung and liver; it is weakly expressed in the spleen; lymph nodes; and thymus. PDCD1LG2 is involved in the costimulatory signal; essential for T-cell proliferation and IFNG production in a PDCD1-independent manner. PDCD1LG2 interaction with PDCD1 inhibits T-cell proliferation by blocking cell cycle progression and cytokine production.