

Purified Anti-Human/Mouse/Rat CD47 Antibody[MIAP410], Functional Grade

catalog number: E-AB-F10160

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Immunogen	Recombinant Human CD47 protein
Host	Mouse
Isotype	Mouse IgG1, κ
Clone	MIAP410
Purification	>98%, Protein A/G purified
Buffer	Sterile PBS, pH 7.2. < 1.0 EU per mg of the antibody as determined by the LAL method.

Applications


Recommended Dilution

FCM	$\leq 0.2 \mu\text{g}$ per million cells in 100 μL volume
Block	Reported in the literature

Data



Rat splenocytes were stained with 0.2 μg Purified Anti-Human/Mouse/Rat CD47 Antibody[MIAP410], Functional Grade (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by FITC-conjugated Goat Anti-Mouse IgG Secondary Antibody.



Rat splenocytes were stained with 0.2 μg Purified Anti-Human/Mouse/Rat CD47 Antibody[MIAP410], Functional Grade (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by FITC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. This preparation contains no preservatives, thus it should be handled under aseptic conditions.
Shipping	Ice bag

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V3.2

Predicted to enable protein binding activity involved in heterotypic cell-cell adhesion and thrombospondin receptor activity. Involved in regulation of nitric oxide biosynthetic process. Acts upstream of or within several processes, including monocyte aggregation; opsonization; and positive regulation of phagocytosis. Located in extracellular exosome. Is integral component of plasma membrane. Is expressed in several structures, including alimentary system; central nervous system; genitourinary system; hemolymphoid system gland; and liver and biliary system. Orthologous to human CD47 (CD47 molecule).

None (Azide-Free, Low Endotoxin) are perfectly suited to be used in culture or in vivo (for nonhuman studies) for functional assays blocking, neutralizing, activation or depletion where the presence of azide may damage cells or exogenous endotoxin may signal or activate cells.

Application References

Yoko Kojima, et al. Nature. 2016 Aug 4;536(7614):86-90.