RHCG Polyclonal Antibody

catalog number: E-AB-53325



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Reactivity	Human	Human	
Immunogen	Synthetic peptide of hun	Synthetic peptide of human RHCG	
Host	Rabbit	Rabbit	
Isotype	IgG	IgG	
Purification	Antigen affinity purificat	Antigen affinity purification	
Conjugation	Unconjugated	Unconjugated	
buffer	Phosphate buffered solu	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.	
Applications	Recommended Dilu	Recommended Dilution	
WB	1:500-1:2000	1:500-1:2000	
IHC	1:25-1:50	1:25-1:50	
Data			
breast tissue lysate Obser	kDa 130	Immunohistochemistry of paraffin-embedded Human brain tissue using RHCG Polyclonal Antibody at dilution of 1:25(×200)	
Preparation & Stor	age		
Storage	Store at -20°C Valid for 12	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.	
Shipping		The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.	

Background

RHCG (Rhesus blood group family type C glycoprotein), also known as RHGK (Rh glycoprotein kidney), PDRC2 (tumorrelated protein DRC2) or SLC42A3, is a multi-pass membrane protein with three potential N-glycosylation sites that belongs to the Rh subfamily of the ammonium transporter family. Expressed in a wide variety of tissues with predominant expression in adult and fetal kidney, RHCg localizes to the apical and/or basolateral cell membrane (depending on the species) and is believed to function as a non-erythroid, bidirectional, electroneutral ammonium transporter.

For Research Use Only