

# PEPD Polyclonal Antibody

Catalog Number:E-AB-52976

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

|                     |                                                        |
|---------------------|--------------------------------------------------------|
| <b>Reactivity</b>   | Human, Mouse, Rat                                      |
| <b>Immunogen</b>    | Fusion protein of human PEPD                           |
| <b>Host</b>         | Rabbit                                                 |
| <b>Isotype</b>      | IgG                                                    |
| <b>Purification</b> | Antigen affinity purification                          |
| <b>Conjugation</b>  | Unconjugated                                           |
| <b>Formulation</b>  | PBS with 0.05% NaN <sub>3</sub> and 40% Glycerol,pH7.4 |

## Applications Recommended Dilution


|              |                |
|--------------|----------------|
| <b>WB</b>    | 1:1000-1:5000  |
| <b>IHC</b>   | 1:50-1:300     |
| <b>ELISA</b> | 1:5000-1:10000 |

## Data




Western blot analysis of HepG2 cell Mouse small intestines tissue lysates using PEPD Polyclonal Antibody at dilution of 1:1000

**Observed Mw:Refer to figures**  
**Calculated Mw:55 kDa**



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using PEPD Polyclonal Antibody at dilution of 1:85(×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using PEPD Polyclonal Antibody at dilution of 1:85(×200)

## Preparation & Storage

**Storage** Store at -20°C. Avoid freeze / thaw cycles.

## Background

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# PEPD Polyclonal Antibody

Catalog Number:E-AB-52976



This gene encodes a member of the peptidase family. The protein forms a homodimer that hydrolyzes dipeptides or tripeptides with C-terminal proline or hydroxyproline residues. The enzyme serves an important role in the recycling of proline, and may be rate limiting for the production of collagen. Mutations in this gene result in prolidase deficiency, which is characterized by the excretion of large amount of di- and tri-peptides containing proline. Multiple transcript variants encoding different isoforms have been found for this gene.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017