

DEAF1 Polyclonal Antibody

catalog number: E-AB-16381

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*


Description

Reactivity	Human
Immunogen	Synthetic peptide of human DEAF1
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications

Applications	Recommended Dilution
IHC	1:50-1:200

Data



Immunohistochemistry of paraffin-embedded Human gastric cancer tissue using DEAF1 Polyclonal Antibody at dilution 1:40

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

Suppressin, also known as DEAF1 (deformed epidermal autoregulatory factor 1), SPN, NUDR (nuclear DEAF-1-related transcriptional regulator), or ZMYND5 (zinc finger MYND domain-containing protein 5), is a transcription factor required for embryonic development. Suppressin contains one SAND domain and one C-terminal MYND-type zinc finger. It interacts with LMO4 and CLIM-2, suggesting that it plays a role mediating cell fate and embryonic pattern formation. Suppressin is expressed in a variety of tissues and localizes to the nucleus. Several isoforms exist due to alternative splicing and, depending on the isoform, Suppressin is secreted in some cell types. Secreted Suppressin can function to inhibit cell proliferation, arresting cells in the G0 or G1 phase. Mutations in the gene encoding Suppressin may result in a growth advantage leading to the development and progression of neoplasia. This suggests that Suppressin is a potential target for cancer therapy.

For Research Use Only