

Human CCL23 Antibody Pair Set

Catalog No.	E-KAB-0184	Applications	ELISA
Synonyms	CCL23; SCYA23; Ckb-8; MIP-3; CKb8; Macrophage inflammatory protein 3		

Kit components & Storage

Title	Specifications	Storage
Human CCL23 Capture Antibody	1 vial, 100 µg	Store at -20°C for one year. Avoid freeze / thaw cycles.
Human CCL23 Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20°C for one year. Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Product Information

Items		Characteristic (E-KAB-0184)	
		Human CCL23 Capture Antibody	Human CCL23 Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Human CCL23 protein	Recombinant Human CCL23 protein
	Swissprot	P55773	
Product details	Reactivity	Human	Human
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4	PBS with 0.04% Proclin 300, 1% protective protein, 50% glycerol, pH 7.4
	Purify	Antigen Affinity	Antigen Affinity
Specificity	Detects Human CCL23 in ELISAs.		

For Research Use Only

Applications

Human CCL23 Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images
ELISA Capture	0.5-4µg/mL	Human CCL23 Capture Antibody	 <p>The graph is a log-log plot. The y-axis is labeled 'Optical Density' and ranges from 0.01 to 10. The x-axis is labeled 'Human MPIF-1/CCL23 concentration(pg/mL)' and ranges from 10 to 10000. Six data points are plotted, showing a clear upward trend. The points are approximately at (30, 0.05), (50, 0.1), (100, 0.2), (200, 0.4), (500, 0.8), and (1000, 1.5).</p>
ELISA Detection	1:1000-1:10000	Human CCL23 Detection Antibody (Biotin)	

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

This gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of the N-terminal cysteine residues of the mature peptide. This chemokine, a member of the CC subfamily, displays chemotactic activity on resting T lymphocytes and monocytes, lower activity on neutrophils and no activity on activated T lymphocytes. The protein is also a strong suppressor of colony formation by a multipotential hematopoietic progenitor cell line.

For Research Use Only