

Recombinant Human SIRPalpha/SIRPA protein (His tag)



Catalog Number:PDEH100330

1 Publications

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|------------------------------------|---|
| Synonyms | Tyrosine-Protein Phosphatase Non-Receptor Type Substrate 1:SHP Substrate 1;SHPS-1;Brain Ig-Like Molecule with Tyrosine-Based Activation Motifs;Bit;CD172 Antigen-Like Family Member A;Inhibitory Receptor SHPS-1;Macrophage Fusion Receptor;MyD-1 Antig |
| Species | Human |
| Expression Host | E.coli |
| Sequence | Glu 31-Tyr 373 |
| Accession | P78324-2 |
| Calculated Molecular Weight | 37.6 kDa |
| Observed molecular weight | 41 kDa |
| Tag | N-His & C-His |

Properties

| | |
|-----------------------|---|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | Please contact us for more information. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Signal Regulatory Protein α (SIRP α) is a monomeric approximately 90 kD type I transmembrane glycoprotein. The 504 amino acid human SIRP α contains two Ig-like C1-type domains and one Ig-like V-type domain. SIRP α can express in various tissues, mainly on brain and myeloid cells, including macrophages, neutrophils, dendritic and Langerhans cells. It also can detect in neurons, smooth muscle and endothelial cells. SIRPA is an immunoglobulin-like cell surface receptor for CD47. SIRP α acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human SIRPalpha/SIRPA protein (His tag)



Catalog Number:**PDEH100330** 1 Publications

the cytosol to the plasma membrane. SIRP α shows adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. SIRP α engagement generally produces a negative regulatory signal; it may mediate negative regulation of phagocytosis, mast cell activation and dendritic cell activation.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017