

Recombinant Human LCN1/VEGP/Lipocalin-1 Protein (His Tag)

Catalog Number: PKSH031078

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species	Human
Source	HEK293 Cells-derived Human LCN1/VEGP/Lipocalin-1 protein Met 1-Asp176, with an C-terminal His
Calculated MW	19.0 kDa
Observed MW	20 kDa
Accession	NP_002288.1
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Lipocalin-1, also known as Von Ebner gland protein, VEG protein, Tear prealbumin, VEGP, Tear lipocalin and LCN1, is a secreted protein which belongs to the calycin superfamily and Lipocalin family. Human Lipocalin-1 / VEGP was originally described as a major protein of human tear fluid, which was thought to be tear specific. Lipocalin-1 / VEGP is identical with lingual von Ebner's gland protein, and is also produced in prostate, nasal mucosa and tracheal mucosa. Homologous proteins have been found in rat, pig and probably dog and horse. Lipocalin-1 / VEGP is an unusual lipocalin member, because of its high promiscuity for relative insoluble lipids and binding characteristics that differ from other members. Lipocalin-1 / VEGP acts as the principal lipid binding protein in tear fluid, a more general physiological function has to be proposed due to its wide distribution and properties. Lipocalin-1 / VEGP would be ideally suited for scavenging of lipophilic, potentially harmful substances and thus might act as a general protection factor of epithelia. Lipocalin-1 / LCN1 could play a role in taste reception. It could be necessary for the concentration and delivery of sapid molecules in the gustatory system. Lipocalin-1 / LCN1 can bind various ligands, with chemical structures ranging from lipids and retinoids to the macrocyclic antibiotic rifampicin and even to microbial siderophores. It exhibits an extremely wide ligand pocket.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017