Recombinant Human Neuropilin-1/NRP1 (C-Avi-6His) Biotinylated

Catalog Number: PKSH033988

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Species	Human
Source	HEK293 Cells-derived Human Neuropilin-1;NRP1 protein Phe22-Lys644, with an C-terminal Avi & His
Calculated MW	72.5 kDa
Observed MW	85-100 kDa
Accession	AAH07533.1
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 ℃. Reconstituted protein solution can be stored at 4-8℃ for 2-7 days. Aliquots of reconstituted samples are stable at < -20℃ for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data

> 95 % as determined by reducing SDS-PAGE.

Background

Neuropilin-1 (Npn-1, previously neuropilin; also CD304) is a 130 - 140 kDa type I transmembrane (TM) glycoprotein that regulates axon guidance and angiogenesis. Two homologues, Neuropilin-1 and Neuropilin-2, are identified. Neuropilin-1 binds to semaphorin 3A, The PLGF-2 isoform of PGF, The VEGF-165 isoform of VEGF and VEGF-B. Coexpression with KDR results in increased VEGF-165 binding to KDR as well as increased chemotaxis. It may regulate VEGF-induced angiogenesis. The soluble isoform 2 binds VEGF-165 and appears to inhibit its binding to cells. NRP1 expression is regulated in EC by tumor necrosis factor-alpha, the transcription factors dHAND and Ets-1, and vascular injury. NRP1 upregulation is positively correlated with the progression of various tumors.

For Research Use Only