



A Reliable Research Partner in Life Science and Medicine

# **Recombinant SerpinD1 Monoclonal Antibody**

catalog number: AN300182P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Reactivity Human

Immunogen Recombinant Human SerpinD1 protein


Host Rabbit
Isotype IgG
Clone A1296
Purification Protein A

Buffer 0.2 µm filtered solution in PBS


Applications Recommended Dilution

**IHC-P** 1:100-1:500

#### Data



Immunohistochemistry of paraffin-embedded human liver using SerpinD1 Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded human hepatoma using SerpinD1 Monoclonal Antibody at dilution of 1:200.

## Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

### **Background**

SerpinD1, also known as heparin cofactor II, is a member of Serpin superfamily of the serine proteinase inhibitors. HCII is a glycoprotein in human plasma that inhibits thrombin and chymotrypsin, and the rate of inhibition of thrombin is rapidly increased by Dermatan sulfate (DS), heparin (H) and glycosaminoglycans(GAG). The stimulatory effect of glycosaminoglycans on the inhibition is mediated, in part, by the N-terminal acidic domain of HCII. Interestingly, a C-terminal His-tagged recombinant HCII exhibits enhanced activity of thrombin inhibition. It has been suggested that HCII plays an unique and important role in vascular homeostasis, and accordingly mutations in this gene or congenital HCII deficiency is potentially associated with thrombosis. HCII specifically inhibits thrombin action at the site of vascular wall injury and HCII-thrombin complexes have been detected in human plasma. HCII protects against thrombin-induced vascular remodeling in both humans and mice and HCII is a predictive biomarker and therapeutic target for atherosclerosis. SerpinD1 also inhibits chymotrypsin, but in a glycosaminoglycan-independent manner.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.2