

S100A4 Polyclonal Antibody

catalog number: E-AB-40459

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;Mouse;Rat
Immunogen	Recombinant Mouse Protein S100-A4 protein
Host	Rabbit
Isotype	IgG
Purification	Antigen Affinity Purification
Buffer	PBS with 0.05% proclin 300, 1% protective protein and 50% glycerol,pH7.4


Applications Recommended Dilution

IHC	1:50-1:200
------------	------------


Data




Immunohistochemistry of paraffin-embedded Human lung using S100A4 Polyclonal Antibody at dilution of 1:50.




Immunohistochemistry of paraffin-embedded Mouse colon using S100A4 Polyclonal Antibody at dilution of 1:50.



Immunohistochemistry of paraffin-embedded Mouse lung using S100A4 Polyclonal Antibody at dilution of 1:50.



Immunohistochemistry of paraffin-embedded Rat colon using S100A4 Polyclonal Antibody at dilution of 1:50.



Immunohistochemistry of paraffin-embedded Rat lung using S100A4 Polyclonal Antibody at dilution of 1:50.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

For Research Use Only

S100A4 is a member of the S100 family of calcium-binding proteins. The S100 family members have been involved in the regulation of a number of cellular processes such as cell cycle progression and differentiation. S100A4 is known to localize to and function in the nucleus, cytoplasm of cells and the extracellular space. S100A4 has also been shown to be associated with tumor growth, motility, invasion, metastasis, angiogenesis, apoptosis and chemoresistance. It is a fibroblast-specific protein associated with mesenchymal cell morphology and motility, is expressed during epithelial-mesenchymal transformations (EMT) in vivo. It is an improved marker for lung fibroblasts that could be useful for investigating the pathogenesis of pulmonary fibrosis. Overexpression of S100A4 is correlated with a worse prognosis in patients with various types of cancer.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.8