

Recombinant Human CCL17 Protein(His Tag)

Catalog Number: GPEH1124

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	E.coli-derived Human CCL17 protein Ala24-Ser94, with an N-terminal His
Calculated MW	7.7 kDa
Observed MW	8 kDa
Accession	Q92583
Bio-activity	Not validated for activity

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human CCL17 proteins, 2µg/lane of
Recombinant Human CCL17 proteins was resolved with
SDS-PAGE under reducing conditions, showing bands at 8
kDa

Background

Chemokines are a family of small chemotactic cytokines, or proteins secreted by cells. Chemokines share the same structure similarities such as small size, and the presence of four cysteine residues in conserved locations in order to form their 3-dimensional shape. Some of the chemokines are considered pro-inflammatory which can be induced to recruit cells of the immune system to a site of infection during an immune response, while others are considered homeostatic and are implied in controlling the migration of cells during normal processes of tissue maintenance and development. There are four members of the chemokine family: C-C kemokines, C kemokines, CXC kemokines and CX3C kemokines. The C-C kemokines have two cysteines nearby the amino terminus. There have been at least 27 distinct members of this subgroup reported for mammals, called C-C chemokine ligands-1 to 28. Chemokine ligand 17 (CCL17), also known as thymus and activation regulated chemokine (TARC), is a small cytokine belonging to the C-C chemokine family. CCL17 is expressed mainly in thymus and transiently in phytohemagglutinin-stimulated peripheral blood mononuclear cells. CCL17 can induce chemotaxis in T cells by binding with the chemokine receptor CCR4.