

Recombinant Human CD74 Protein (His Tag)

Catalog Number: PKSH031262

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human CD74 protein Gln 73-Met 232, with an N-terminal His
Calculated MW	20.4 kDa
Accession	NP_004346.1
Bio-activity	Immobilized human CD74 at 5 µg/ml (100 µl/well) can bind biotinylated human CTSL1 with a linear range of 3. 2-400 ng/ml.

Properties

Purity	> 80 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 80 % as determined by reducing SDS-PAGE.

Background

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD74, also known as HLA class2 histocompatibility antigen gamma chain and HLA-DR antigens-associated invariant chain, is a polypeptide involved in the formation and transport of MHC class2 protein. CD74 is expressed by B cells, macrophages, and Reed-Sternberg cells. When MHC class 2 protein was in the rough ER, its peptide-binding cleft was blocked by CD74 to prevent it from interacting with the endogenous peptides. CD74 also serves to facilitate MHC class2's export from ER.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017