

Recombinant Human Follistatin 288/FST Protein

Catalog Number: PKSH033674

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human Follistatin 288;FST protein Gly30-Asn317, with an C-terminal His
Calculated MW	32.4 kDa
Observed MW	33-42 kDa
Accession	P19883
Bio-activity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Follistatin 288 is a secreted glycoprotein that was first identified as a follicle-stimulating hormone inhibiting substance in ovarian follicular fluid. Human follistatin 288 cDNA encodes a 317 amino acid (aa) protein with a 29aa signal sequence, and a 288 aa mature region. Follistatin shows the highest affinity for activins due to its extended configuration. Genetic deletion of follistatin in mice, or expression of only the Follistatin form, is perinatally lethal due to defects of lung, skin and musculoskeletal system. Follistatins also regulate hematopoietic stem cell adhesion to fibronectin via FS2.

For Research Use Only