

Human AGER Antibody Pair Set

Catalog No.	E-KAB-0217	Applications	ELISA
Synonyms	AGER, RAGE, Receptor for Advanced Glycation End products,		

Kit components & Storage

Title	Specifications	Storage
Human AGER Capture Antibody	1 vial, 100 µg	Store at -20°C for one year. Avoid freeze / thaw cycles.
Human AGER Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20°C for one year. Avoid freeze / thaw cycles.


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0217)	
		Human AGER Capture Antibody	Human AGER Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Human AGER protein	Recombinant Human AGER protein
	Swissprot	Q15109	
Product details	Reactivity	Human	Human
	Host	Mouse	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4	PBS with 0.04% Proclin 300, 1% protective protein, 50% glycerol, pH 7.4
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Human AGER in ELISAs.	

Applications

Human AGER Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images																
ELISA Capture	0.5-4µg/mL	Human AGER Capture Antibody	 <p>The graph is a log-log plot. The x-axis is labeled 'Human AGER concentration(pg/mL)' and ranges from 10 to 100,000. The y-axis is labeled 'Optical Density' and ranges from 0.01 to 10. The data points form a straight line with a positive slope, indicating a power-law relationship between concentration and optical density.</p> <table border="1"> <caption>Approximate data points from the standard curve</caption> <thead> <tr> <th>Human AGER concentration (pg/mL)</th> <th>Optical Density</th> </tr> </thead> <tbody> <tr> <td>100</td> <td>0.05</td> </tr> <tr> <td>200</td> <td>0.1</td> </tr> <tr> <td>500</td> <td>0.25</td> </tr> <tr> <td>1000</td> <td>0.5</td> </tr> <tr> <td>2000</td> <td>1.0</td> </tr> <tr> <td>5000</td> <td>2.5</td> </tr> <tr> <td>10000</td> <td>5.0</td> </tr> </tbody> </table>	Human AGER concentration (pg/mL)	Optical Density	100	0.05	200	0.1	500	0.25	1000	0.5	2000	1.0	5000	2.5	10000	5.0
Human AGER concentration (pg/mL)	Optical Density																		
100	0.05																		
200	0.1																		
500	0.25																		
1000	0.5																		
2000	1.0																		
5000	2.5																		
10000	5.0																		
ELISA Detection	1:1000-1:10000	Human AGER Detection Antibody (Biotin)																	

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

The advanced glycosylation end product (AGE) receptor encoded by this gene is a member of the immunoglobulin superfamily of cell surface receptors. It is a multiligand receptor, and besides AGE, interacts with other molecules implicated in homeostasis, development, and inflammation, and certain diseases, such as diabetes and Alzheimer's disease. Many alternatively spliced transcript variants encoding different isoforms, as well as non-protein-coding variants, have been described for this gene.