

Mouse IL-17A Antibody Pair Set

Catalog No.	E-KAB-0082	Applications	ELISA
Synonyms	CTLA8, CTLA-8, CTLA8cytotoxic T-lymphocyte-associated serine esterase 8, Cytotoxic T-lymphocyte-associated antigen 8, IL17, IL-17, IL17A, IL-17A, IL-17Acytotoxic T-lymphocyte-associated protein 8, IL-17CTLA-8		

Kit components & Storage

Title	Specifications	Storage
Mouse IL-17A Capture Antibody	1 vial, 100 µg	Store at -20°C for one year. Avoid freeze / thaw cycles.
Mouse IL-17A Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20°C for one year. Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Product Information

Items		Characteristic (E-KAB-0082)	
		Mouse IL-17A Capture Antibody	Mouse IL-17A Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Mouse IL-17A protein	Recombinant Mouse IL-17A protein
	Swissprot	Q62386	
Product details	Reactivity	Mouse	Mouse
	Host	Goat	Rat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4	PBS with 0.04% Proclin 300, 1% protective protein, 50% glycerol, pH 7.4
	Purify	Antigen Affinity	Protein A or G
	Specificity	Detects Mouse IL-17A in ELISAs.	

For Research Use Only

Applications

Mouse IL-17A Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images										
ELISA Capture	0.5-4μg/mL	Mouse IL-17A Capture Antibody	 <table><caption>Approximate data points from the standard curve</caption><thead><tr><th>Mouse IL-17A concentration (pg/mL)</th><th>Optical Density</th></tr></thead><tbody><tr><td>10</td><td>0.05</td></tr><tr><td>100</td><td>0.2</td></tr><tr><td>1000</td><td>1.0</td></tr><tr><td>10000</td><td>5.0</td></tr></tbody></table>	Mouse IL-17A concentration (pg/mL)	Optical Density	10	0.05	100	0.2	1000	1.0	10000	5.0
Mouse IL-17A concentration (pg/mL)	Optical Density												
10	0.05												
100	0.2												
1000	1.0												
10000	5.0												
ELISA Detection	1:1000-1:10000	Mouse IL-17A Detection Antibody (Biotin)											

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

IL17a is a pro-inflammatory cytokine as a member of the interleukin-17 family. It plays a central role in host defense against diverse pathogens. It is produced by activated T-cells and certain cell types of innate immune system. The active protein functions as either a homodimer with other interleukin-17 family members and signals through the interleukin-17 receptor to induce inflammatory cytokine production. Aberrant expression of this gene is associated with autoinflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis.

For Research Use Only