

A Reliable Research Partner in Life Science and Medicine

# PE Anti-Human CD195 Antibody[RoAb13]

Catalog Number: E-AB-F1418D

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human Host Mouse

**Isotype** Mouse IgG2a, κ

Clone No. RoAb13

Isotype Control PE Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802D]

Conjugation PE

Conjugation Information PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green

(561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42

nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.


### Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

#### Data



Staining of normal human peripheral blood cells with APC Anti-Human CD3 Antibody and PE Anti-Human CD195 Antibody[RoAb13] (left) or PE Mouse IgG2a,  $\kappa$  Isotype Control (right). Cells in the lymphocytes gate were used for analysis.

### **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Web: www.elabscience.cn

Shipping Ice bag

# **Antigen Information**

 Uniprot ID
 P51681

 Gene ID
 1234

### For Research Use Only



## Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

### **Background**

CD195, also known as CCR5, is a 45 kD G protein-coupled seven transmembrane C C-chemokine receptor. It binds to MIP-1 $\alpha$ , MIP-1 $\beta$ , and RANTES and is expressed on a subset of T cells and monocytes. CCR5 mediates an intracellular signal thought to induce cell differentiation and proliferation. CCR5 has also been shown to act as a coreceptor for R5 HIV-1 cell entry; modification of CCR5 by sulfation contributes to the efficiency of HIV-1 entry. Studies have shown CCR5 to play a role in a variety of other human diseases, ranging from infectious and inflammatory diseases to cancer.

Web: www.elabscience.cn