

A Reliable Research Partner in Life Science and Medicine

IL-17 RA/IL-17 R Polyclonal Antibody(Capture/Detector)

catalog number: AN000470P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

т					٠.		
- 11	es	O I	ra	m	ш	\mathbf{a}	т

Reactivity Human

Immunogen Recombinant Human IL-17 RA/IL-17 R RA protein expressed by Mammalian

Host Rabbit Isotype Rabbit IgG

Purification Antigen Affinity Purification


Conjugation Unconjugated

Buffer Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

Applications Recommended Dilution

ELISA Capture 2-8 μg/mL ELISA Detector 0.1-0.4 μg/mL

Data



Sandwich ELISA-Recombinant Human IL-17 RA/IL-17 R RA protein standard curve.Background subtracted standard curve using IL-17 RA/IL-17 R antibody(AN000470P)

(Capture), IL-17 RA/IL-17 R antibody (AN000470P)

(Detector) in sandwich ELISA. The reference range value for Recombinant Human IL-17 RA/IL-17 R RA protein is 0.31-

20 pg/mL.

20 pg/m

Preparation & Storage

Storage Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

Interleukin 17A (IL17A) is a proinflammatory cytokine secreted by activated T-lymphocytes. It is a potent inducer of the maturation of CD34-positive hematopoietic precursors into neutrophils. The transmembrane protein encoded by this gene (interleukin 17A receptor; IL17RA) is a ubiquitous type I membrane glycoprotein that binds with low affinity to interleukin 17A. Interleukin 17A and its receptor play a pathogenic role in many inflammatory and autoimmune diseases such as rheumatoid arthritis. Like other cytokine receptors, this receptor likely has a multimeric structure. Alternative splicing results in multiple transcript variants encoding different isoforms.

Web: www.elabscience.cn

For Research Use Only