PLA2G16 Polyclonal Antibody

catalog number: **E-AB-53452**


Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description


Reactivity	Human;Mouse;Rat
Immunogen	Synthetic peptide of human PLA2G16
Host	Rabbit
Isotype	Rabbit IgG
Purification	Antigen affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications	Recommended Dilution
IHC	1:30-1:150

Data



Immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using PLA2G16 Polyclonal Antibody at dilution of 1:40(×200)



Immunohistochemistry of paraffin-embedded Human tonsil tissue using PLA2G16 Polyclonal Antibody at dilution of 1:40(×200)

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

Background

Secretory phopholipase A2 (PLA2) enzymes cleave an acyl ester bond in the sn-2 position of glycerophospholipids. These extracellular proteins have a high disulfide bond content, low molecular mass (14 kDa), and require mM levels of Ca2+ for catalysis. They play a crucial role in the generation of arachidonates and eicosanoids, and have a number of biological actions including immunological responses, inflammation, cellular proliferation, vasoconstriction, and bronchioconstriction. Exhibits PLA1/2 activity, catalyzing the calcium-independent hydrolysis of acyl groups in various phosphatidylcholines (PC) and phosphatidylethanolamine (PE). For most substrates, PLA1 activity is much higher than PLA2 activity. Specifically catalyzes the release of fatty acids from phospholipids in adipose tissue (By similarity). N- and O-acylation activity is hardly detectable. Might decrease protein phosphatase 2A (PP2A) activity.

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com