

Recombinant Mouse CD150/SLAM Protein (His Tag)

Catalog Number: PKSM041149

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse CD150/SLAM protein Thr25-Pro242 , with an C-terminal His
Calculated MW	25.2 kDa
Observed MW	40-60 kDa
Accession	Q9QUM4
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V3.6

Signaling lymphocyte activation molecule (SLAM), is a self-ligand glycoprotein which exists not only found on the surface of activated and memory T cells, but also on the surface of activated B cells, dendritic cells, and macrophages. SLAM consists of a extracellular domain (ECD) with two Ig-like domains,transmembrane segment, and cytoplasmic domain with three immunoreceptor tyrosine switch motifs (ITSM). SLAM is thought to play an important role in adhesion between T cells and APCs and has been shown to act as a coreceptor in TCR-dependent responses. SLAM, together with CD46, is one of the two receptors for measles virus. SLAM is a cell surface receptor that, like the B cell receptor, CD40, and CD95, can transmit positive or negative signals. SLAM can associate with the SH2-containing inositol phosphatase (SHIP), the SH2-containing protein tyrosine phosphatase (SHP-2), and the adaptor protein SH2 domain protein 1A. It's upregulated on activated B cells and CD4+ and CD8+ T cells, but downregulated on Th2 polarized cells. Also, it can Inhibits antigen receptor-mediated production of IFN-gamma, but not IL-2, in CD4-/CD8- T-cells