

**Flt-3 Ligand/FL/FLG3L/FLT3L, Mouse, Recombinant****Cat. No. : GPCK356****产品信息**

物种	Mouse
表达宿主	E.coli
序列信息	MGTPDCYFSHSPISSNFKVKFRELTDLHKDYPVTVAVNLQDEKHCKALWSLFLAQRWIEQLK TVAGSKMQTLLEDVNTEIHFVTSCTFQPLPECLRFVQTNISHLLKDTCTQLLALKPCIGKACQN FSRCLEVQCQPDSSLLPPRSPIALEATELPEPRPR with polyhistidine tag at the C-terminus.
检索号	P49772
标签	His-tag at the C-terminus
分子量	19.3 kDa
有效期	12 months
生物活性	Measure by its ability to induce proliferation in BaF3 cells transfected with mouse Flt-3. The ED50 for this effect is < 2 ng/mL.

**产品特性**

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 μm filtered solution containing 1 × PBS, pH 7.4.
复溶	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

**背景介绍**

Fms-related tyrosine kinase 3 ligand (FLT3LG) is a protein which in humans is encoded by the FLT3LG gene. FLT3 ligand is a receptor for the fl cytokine has a tyrosine-protein kinase activity & a growth factor that regulates proliferation of early hematopoietic cells. Flt3-Ligand synergizes with other CSFs and interleukins to induce growth and differentiation.

