

Recombinant Human/Rhesus HER4/ErbB4 Protein (Fc Tag)

Catalog Number: PKSH031650

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human HER4/ErbB4 protein Met 1-Arg649, with an C-terminal hFc
Mol_Mass	96.6 kDa
Accession	NP_005226.1
Bio-activity	1. Measured by its ability to bind biotinylated human Fc-NRG1 (isoform Beta1) in a functional ELISA. 2. Measured by its ability to bind biotinylated human NRG1 (isoform Beta1) in a functional ELISA. 3. Measured by its ability to bind biotinylated human NRG1 (aa 2-246)-Fc in a functional ELISA.

Properties

Purity	> 85 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Reconstitution	Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information.

Data



> 85 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V3.4

ERBB4 is a single-pass type I membrane protein with multiple cysteine rich domains; a transmembrane domain; a tyrosine kinase domain; a phosphatidylinositol-3 kinase binding site and a PDZ domain binding motif. ERBB4 is expressed at highest levels in brain; heart; kidney; in addition to skeletal muscle; parathyroid; cerebellum; pituitary; spleen; testis and breast. And lower levels in thymus; lung; salivary gland; and pancreas. It specifically binds to and is activated by neuregulins; NRG-2; NRG-3; heparin-binding EGF-like growth factor; betacellulin and NTAK. ERBB4 also can be activated by other factors and induces a variety of cellular responses including mitogenesis and differentiation. ERBB4 regulates development of the heart; the central nervous system and the mammary gland; gene transcription; cell proliferation; differentiation; migration and apoptosis. It is required for normal cardiac muscle differentiation during embryonic development; and for postnatal cardiomyocyte proliferation. ERBB4 also play a role on the normal development of the embryonic central nervous system; especially for normal neural crest cell migration and normal axon guidance. It is required for mammary gland differentiation; induction of milk proteins and lactation.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017