

## Recombinant Human Nicastrin/NCSTN Protein (His Tag)

**Catalog Number:** PKSH031225

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human Nicastrin/NCSTN protein Met 1-Glu 669, with an C-terminal His
<b>Calculated MW</b>	72.4 kDa
<b>Observed MW</b>	110-120 kDa
<b>Accession</b>	NP_056146.1
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 93 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 93 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

Nicastrin (NCST, or NCT), a single-pass membrane glycoprotein that harbors a large extracellular domain, is an essential component of the gamma-secretase complex. Several lines of evidence indicate that the members of these complexes could also contribute to the control of cell death. NCT controls cell death via phosphoinositide 3-kinase/Akt and p53-dependent pathways and that this function remains independent of the activity and molecular integrity of the gamma-secretase complexes. Increasing evidences have shown that Nicastrin/NCSTN plays a crucial role in gamma-cleavage of the amyloid precursor protein (APP). The glycoprotein Nicastrin is an essential component of the gamma-secretase complex, a high molecular weight complex which also contains the presenilin proteins, Aph-1 and Pen-2. The gamma-secretase complex is not only involved in APP processing but also in the processing of an increasing number of other type I integral membrane proteins. As the largest subunit of the gamma-secretase complex, Nicastrin plays a crucial role in its activation. Inhibition of NCSTN demonstrated an altered gamma-cleavage activity, suggesting its potential implication in developing Alzheimer's disease (AD). In addition, Nicastrin can function to maintain epithelial to mesenchymal transition during breast cancer progression. Anti-nicastrin polyclonal and monoclonal antibodies were able to decrease notch1 and vimentin expression and reduced the invasive capacity of breast cancer cells in vitro.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017