



A Reliable Research Partner in Life Science and Medicine

Recombinant Human CXCL9 protein(His Tag)

Catalog Number: PKSH034195

Note: Centrifuge before opening to ensure complete recovery of vial contents.

\mathbf{r}					٠.			
H))	es	C	r٦	n	tı	n	m	

Species Human

Source E.coli-derived Human CXCL9 protein Thr 23-Thr 125, with an N-terminal His

 Calculated MW
 12.5 kDa

 Observed MW
 18 kDa

 Accession
 Q07325

Bio-activity Measure by its ability to chemoattract BaF3 cells transfected with mouse CXCR3. The

ED₅₀ for this effect is $< 0.5 \mu g/mL$.

Properties

Purity > 98 % as determined by reducing SDS-PAGE.

Endotoxin < 0.1 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4.

Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Background

Chemokine (C-X-C Motif) Ligand 9 (CXCL9) belongs to the intercrine alpha (chemokine CXC) family. It is secreted by interferon stimulated monocytes, macrophages and endothelial cells, which elicits chemotactic functions by interacting with the chemokine receptor CXCR3. CXCL9 acts as a Th1 (type 1 helper T) cell chemoattractant and plays a role in the growth, activation and movement of cells associated with immune and inflammatory responses, and in tumour growth inhibition. It is closely related to two other CXC chemokines called CXCL10 and CXCL11, whose genes are located near the gene for CXCL9 on human chromosome 4.