# Purified Anti-Human CD194/CCR4 Antibody

**catalog number:** **E-AB-F1366A**


**Note:** *Centrifuge before opening to ensure complete recovery of vial contents.*

## Description

| | |
|---|---|
| **Reactivity** | Human |
| **Immunogen** | Recombinant Human CD194 protein |
| **Host** | Mouse |
| **Isotype** | Mouse IgG1, κ |
| **Clone** | L291H4 |
| **Purification** | >98%, Protein A/G purified |
| **Conjugation** | Unconjugated |
| **Buffer** | Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling. |

| Applications | Recommended Dilution |
|---|---|
| **FCM** | ≤ 0.2 μg per million cells in 100 μL volume |

## Data



Human peripheral blood lymphocytes cell were stained with 0.2 μg Purified Anti-Human CD194/CCR4 Antibody[L291H4] (Right) and 0.2 μg Mouse IgG1, κ Isotype Control (Left), followed by Elab Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD3 PE-conjugated Monoclonal Antibody.

## Preparation & Storage

| | |
|---|---|
| **Storage** | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles. |
| **Shipping** | Ice bag |

## Background

High affinity receptor for the C-C type chemokines CCL17/TARC, CCL22/MDC and CKLF isoform 1/CKLF1. The activity of this receptor is mediated by G(i) proteins which activate a phosphatidylinositol-calcium second messenger system. Can function as a chemoattractant homing receptor on circulating memory lymphocytes and as a coreceptor for some primary HIV-2 isolates. In the CNS, could mediate hippocampal-neuron survival.


Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com