
Elabscience®

Purified Anti-Human CD194 Antibody[L291H4]

catalog number: E-AB-F1366A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|--|--|
| Reactivity | Human |
| Immunogen | Recombinant Human CD194 protein |
| Host | Mouse |
| Isotype | Mouse IgGl, κ |
| Clone | L291H4 |
| Purification | >98%, Protein A/G purified |
| Buffer | Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling. |
| Applications | Recommended Dilution |
| FCM | $2 \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$ |
| Data | |
| and Tanana (120- 14-14-14-14-14-14-14-14-14-14-14-14-14-1 | |



Human peripheral blood lymphocytes cell were stained with 0.2 µg Purified Anti-Human CD194 Antibody[L291H4]

(Right) and 0.2 μ g Mouse IgG1, κ Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Mouse

All a second and a

IgG Secondary Antibody, then anti-Human CD3 PEconjugated Monoclonal Antibody

| conjugated Monocional Antibody. | |
|---------------------------------|---|
| Preparation & Storage | |
| Storage | Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / |
| | thaw cycles. |
| Shipping | Ice bag |
| Background | |

High affinity receptor for the C-C type chemokines CCL17/TARC, CCL22/MDC and CKLF isoform 1/CKLF1. The activity of this receptor is mediated by G(i) proteins which activate a phosphatidylinositol-calcium second messenger system. Can function as a chemoattractant homing receptor on circulating memory lymphocytes and as a coreceptor for some primary HIV-2 isolates. In the CNS, could mediate hippocampal-neuron survival.

For Research Use Only