## **Elabscience Biotechnology Co., Ltd.**



A Reliable Research Partner in Life Science and Medicine

# Purified Anti-Human CD11a Antibody[R7-1]

Catalog Number: GF1211A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Reactivity Human

Immunogen Recombinant Human CD11A protein

**Host** Mouse

**Isotype** Mouse IgG1,κ

Clone R7-1

**Purification** >98%, Protein A/G purified


**Conjugation** Unconjugated

**Buffer** PBS, pH 7.2. Contains 0.05% proclin 300.

### Applications Recommended Dilution

FCM  $2 \mu g/mL(0.5 \times 10^6 - 1 \times 10^6 \text{ cells})$ 

#### Data



Human peripheral blood granulocytes cell were stained with 0.2  $\mu$ g Purified Anti-Human CD11a Antibody[R7-1] (Right) and 0.2  $\mu$ g Mouse IgG1,  $\kappa$  Isotype Control (Left), followed by

Elab Fluor® 647-conjugated Goat Anti-Mouse IgG Secondary Antibody.

### **Preparation & Storage**

Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid

freeze / thaw cycles.

Shipping Ice bag

#### **Background**

CD11a is a 170-180 kD type I transmembrane glycoprotein also known as LFA-1 $\alpha$  chain and integrin  $\alpha$ L subunit. CD11a non-covalently associates with integrin  $\beta$ 2 (CD18) to form LFA-1. It is expressed on all leukocytes, including B and T lymphocytes, monocytes, macrophages, neutrophils, basophils and eosinophils. It is absent on non-hematopoietic tissues and platelets. CD11a plays a central role in leukocyte cell-cell interactions and is important in lymphocyte costimulation. CD11a/CD18 binds to ICAM-1 (CD54), ICAM-2 (CD102), and ICAM-3 (CD50).