



A Reliable Research Partner in Life Science and Medicine

Biotin Anti-Human CD38 Antibody[HIT2]

Catalog Number: E-AB-F1058B

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human
Host Mouse

Isotype Mouse IgG1, κ

Clone No. HIT2

Isotype Control Biotin Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09793B]

Conjugation Biotin

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein

protectant.

Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. For flow

cytometric staining, the suggested use of this reagent is \leq 1.0 μ g per 10⁶ cells in 100 μ L volume or 100 μ L of whole blood. It is recommended that the reagent be titrated for

optimal performance for each application.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names 2'-phospho-ADP-ribosyl cyclase;2'-phospho-cyclic-ADP-ribose transferase;ADP-ribosyl

cyclase 1;ADP-ribosyl cyclase/cyclic ADP-ribose hydrolase 1;ADPRC 1;CD38;Cyclic

ADP-ribose hydrolase 1;T10;cADPr hydrolase 1

 Uniprot ID
 P28907

 Gene ID
 952

Background CD38 is a 45 kD type II transmembrane glycoprotein also known as T10. It is an ADP-

ribosyl hydrolase expressed at variable levels on hematopoietic cells and in some non-

hematopoietic tissues (such as brain, muscles, and kidney). In humans, it is

expressed at high levels on plasma cells and activated T and B cells. By functioning as both a cyclase and a hydrolase, CD38 mediates lymphocyte activation, adhesion, and

the metabolism of cADPR and NAADP. CD31 is the ligand of CD38.