Recombinant Human Interleukin-16/IL-16 Protein

Catalog Number: PKSH032618



Description			
Species	Human		
Mol_Mass	13.4 kDa		
Accession	AAC12732.1		
Bio-activity	Not validated for activity		
Properties			
Purity	> 95 % as determined by reducing SDS-PAGE.		
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.		
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80		
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.		
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Formulation	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.0.		
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants		
	before lyophilization.		
	Please refer to the specific buffer information in the printed manual.		
Reconstitution	Please refer to the printed manual for detailed information.		

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Data		
	kDa	MK
	120 90	
	60	- abscien
	40	
	30	e medience
	20	and Bar
		had she
	14	

> 95 % as determined by reducing SDS-PAGE.

Background

Interleukin-16 (IL-16) is a CD8+ T cell-derived cytokine that induces chemotaxis of CD4+ T cells and CD4+ monocytes and eosinophils. Analysis by gel filtration suggests that, under physiological conditions, human IL-16 exists predominantly as a noncovalently linked multimer, but that some IL-16 may exist as a monomer. However, only the multimeric form appears to possess chemotactic activity, suggesting that receptor cross-linking may be required for activity. IL-16 also induces expression of IL-2 receptor (IL-2R) and MHC class II molecules on CD4+ T cells. Human and murine IL-16 show significant cross-species reactivity.

For Research Use Only