

Biotin Anti-Human CD64 Antibody[10.1]

Catalog Number: E-AB-F1082B

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, κ
Clone No.	10.1
Isotype Control	Biotin Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09793B]
Conjugation	Biotin
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \mu\text{g}$ per 10^6 cells in 100 μL volume or 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
------------	---

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	CD64;CD64A/B/C;FCGR1A/B/C;Fc fragment of IgG high affinity Ia/b/c receptor;Fc gamma RI;IGFR 1
Uniprot ID	P12314
Gene ID	2209
Background	CD64 is a 72 kD single chain type I glycoprotein also known as FcγRI and FcR I. CD64 is a member of the immunoglobulin superfamily and is expressed on monocytes/macrophages, dendritic cells, and activated granulocytes. The expression can be upregulated by IFN-γ stimulation. CD64 binds IgG immune complex. It plays a role in antigen capture, phagocytosis of IgG/antigen complexes, and antibody-dependent cellular cytotoxicity (ADCC).

For Research Use Only