

## DNAJC17 Polyclonal Antibody

**catalog number: E-AB-92184**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Reactivity</b>	Human;Mouse;Rat
<b>Immunogen</b>	Recombinant fusion protein of human DNAJC17
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Purification</b>	Affinity purification
<b>Buffer</b>	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

### Applications


<b>WB</b>	1:1000-1:2000
<b>IHC</b>	1:50-1:200

### Data




Western blot analysis of extracts of various cell lines using DNAJC17 Polyclonal Antibody at 1:1000 dilution.

**Observed-MW:35 kDa**  
**Calculated-MW:35 kDa**




Immunohistochemistry of paraffin-embedded human esophageal cancer using DNAJC17 Polyclonal Antibody at dilution of 1:20 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Western blot analysis of extracts of various cell lines using DNAJC17 Polyclonal Antibody at 1:1000 dilution.

**Observed-MW:35 kDa**  
**Calculated-MW:35 kDa**



Immunohistochemistry of paraffin-embedded mouse spinal cord using DNAJC17 Polyclonal Antibody at dilution of 1:20 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

### Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

### Background

May negatively affect PAX8-induced thyroglobulin/TG transcription.

### For Research Use Only